

LETTER • OPEN ACCESS

Evidencing decadal and interdecadal hydroclimatic variability over the Central Andes

To cite this article: Hans Segura *et al* 2016 *Environ. Res. Lett.* **11** 094016

View the [article online](#) for updates and enhancements.

Related content

- [Facing unprecedented drying of the Central Andes? Precipitation variability over the period AD 1000–2100](#)
Raphael Neukom, Mario Rohrer, Pierluigi Calanca *et al.*
- [Is anthropogenic sea level fingerprint already detectable in the Pacific Ocean?](#)
H Palanisamy, B Meyssignac, A Cazenave *et al.*
- [A dipole pattern in the Indian and Pacific oceans and its relationship with the East Asian summer monsoon](#)
Jiayu Zheng, Jianping Li and Juan Feng

Recent citations

- [Linking atmospheric circulation patterns with hydro-geomorphic disasters in Peru](#)
C. Rodríguez-Morata *et al*
- [Rainfall along the coast of Peru during strong El Niño events](#)
Janeet Sanabria *et al*
- [Understanding the influence of orography on the precipitation diurnal cycle and the associated atmospheric processes in the central Andes](#)
C. Junquas *et al*

Environmental Research Letters



LETTER

Evidencing decadal and interdecadal hydroclimatic variability over the Central Andes

OPEN ACCESS

RECEIVED

15 February 2016

REVISED

31 July 2016

ACCEPTED FOR PUBLICATION

8 August 2016

PUBLISHED

14 September 2016

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Hans Segura¹, Jhan Carlo Espinoza¹, Clementine Junquas² and Ken Takahashi¹¹ Instituto Geofísico del Perú (IGP), Lima, Peru² IRD/UGA/CNRS/G-INP, LTRE UMR 5564, Grenoble, FranceE-mail: jhan-carlo.espinoza@igp.gob.pe**Keywords:** decadal variability, Central Andes, South America, climate variability, hydrology, El NiñoSupplementary material for this article is available [online](#)


Abstract

In this study we identified a significant low frequency variability (8 to 20 years) that characterizes the hydroclimatology over the Central Andes. Decadal–interdecadal variability is related to the central-western Pacific Ocean ($R^2 = 0.50$) and the zonal wind at 200 hPa above the Central Andes ($R^2 = 0.66$). These two oceanic–atmospheric variables have a dominant decadal–interdecadal variability, and there is a strong relationship between them at a low frequency time scale ($R^2 = 0.66$). During warming decades in the central-western Pacific Ocean, westerlies are intensified at 200 hPa above the Central Andes, which produce decadal periods of hydrological deficit over this region. In contrast, when the central-western Pacific Ocean is cooler than usual, easterly anomalies prevail over the Central Andes, which are associated with decades of positive hydrological anomalies over this region. Our results indicate that impacts of El Niño on hydrology over the Central Andes could be influenced by the low frequency variability documented in this study.

Introduction

Precipitation in the Central Andes is characterized by a strong spatial contrast due to its complex topographic gradient (Garreaud *et al* 2003, Vuille and Keimig 2004) (figures 1(a) and (b)). In this region, the Andes form a natural barrier for warm and wet easterly winds from the central Amazon Basin and the Atlantic Ocean (Melice and Roucou 1998, Garreaud 1999, Garreaud *et al* 2003, Vuille *et al* 2003, Garreaud 2009, Insel *et al* 2010), and cold, dry winds from the subtropical Pacific Ocean (Garreaud 1999, Houston and Hartley 2003). While the eastern part of the Andes is characterized by atmospheric instability, deep convection and a huge amount of rainfall (Garreaud 1999, Garreaud 2000, Vuille *et al* 2000, Garreaud *et al* 2003, Vuille *et al* 2003, Vuille and Keimig 2004, Espinoza *et al* 2015), the Western Andes are characterized by great atmospheric stability due to large-scale subsidence and a cool subtropical east Pacific Ocean (Garreaud 1999, Houston and Hartley 2003, Rutllant *et al* 2003, Garreaud 2009).

Precipitation over the Central Andes is the principal water source for the dry southern coast of Peru, the Altiplano region and the upper part of the Amazon. In these regions, rainfall shows a strong seasonality; indeed, more than 50% of the precipitation falls during the austral summer—December to March—(Vuille *et al* 2000, Garreaud 2000, Garreaud *et al* 2003, Vuille and Keimig 2004, Knüsel *et al* 2005). This seasonality is related to the presence of the Bolivian High (BH) in the upper troposphere (Lenters and Cook 1997, Vuille 1999, Vuille *et al* 2000, Garreaud *et al* 2003, Garreaud 2009), which is formed by convection in the central Amazon (Silva Dias *et al* 1983, Lenters and Cook 1997). The BH is associated with an anticyclonic and divergent movement of winds at 200 hPa, which allows zonal winds to carry moister air from the inner part of the continent (Vuille 1999, Garreaud *et al* 2003). A significant correlation has therefore been documented between rainfall in the Altiplano and easterly winds at 200 hPa over the Central Andes, which have been observed from synoptic to interannual



time scales (Garreaud *et al* 2003, Minvielle and Garreaud 2011).

Previous studies have noted that the Tropical Pacific plays a relevant role in modulating rainfall in the Central Andes on an interannual time scale (Aceituno and Garreaud 1995, Ronchail 1995, Vuille 1999, Vuille *et al* 2000, Garreaud and Aceituno 2001, Vuille *et al* 2003, Garreaud *et al* 2003, Lagos *et al* 2008, Lavado-Casimiro and Espinoza 2014). Lavado-Casimiro and Espinoza (2014) found that central El Niño events are distinguished by a precipitation decrease in the Andes, especially in the Central Andes. High (low) values of the Southern Oscillation Index are also associated with easterly (westerly) anomalies at 200 hPa over this region, which are characteristic of wet (dry) years (Vuille 1999, Garreaud and Aceituno 2001). El Niño 1982–1983 and 1997–1998 had different impacts on precipitation and hydrology in the Central Andes; for example, El Niño 1982–1983 was characterized by strong deficit of precipitation throughout the Central Andes, while El Niño 1997–1998 did not cause the same effects (Sperling *et al* 2008).

In the western portion of the Central Andes, a decadal–interdecadal rainfall oscillation was detected and was statistically related to sea surface temperature (SST) in the central and eastern tropical Pacific and the

South Pacific Convergence Zone (Vuille *et al* 2000). Seiler *et al* (2013) suggested that extreme warm and dry events in the Bolivian highlands are more frequent during positive phase of the Pacific Decadal Oscillation (PDO), while a negative phase of the PDO enhanced floods in the Bolivian lowlands. Although a few studies have explored low frequency climate variability over the central and western part of the Altiplano, decadal/interdecadal rainfall variability over all the Central Andes remains poorly documented. There is a clear need for in-depth analysis of low frequency hydroclimatic variability over this region and its relationship to oceanic and atmospheric conditions.

In this study we use a hydroclimatic dataset as a proxy for precipitation in the Central Andes to analyze its low frequency variability. We use discharge data from rivers that drain into the Pacific (western Central Andes) and into the Amazon basin (eastern Central Andes), as well as the water level in Lake Titicaca, which synthesizes rainfall variability in the Altiplano (Aceituno and Garreaud 1995). In addition, the causes of low frequency hydroclimatic variability over the Central Andes are analyzed and discussed in relation to low frequency variability of global SST and atmospheric circulation.

Table 1. Name, geographic location, temporal range, hydrological system and type of hydrological data for each station.

Name of station	Location	Temporal range (years)	Hydrological systems	Hydrological data
Aguas Calientes	70.12 °W; 17.85 °S	1969–2005	Pacific	Discharge ($\text{m}^3 \text{s}^{-1}$)
La Tranca	70.48 °W; 17.73 °S	1969–2005	Pacific	Discharge ($\text{m}^3 \text{s}^{-1}$)
La Pascana	71.70 °W; 17.03 °S	1956–1999	Pacific	Discharge ($\text{m}^3 \text{s}^{-1}$)
Puente Santa Rosa	71.63 °W; 16.98 °S	1990–2014	Pacific	Discharge ($\text{m}^3 \text{s}^{-1}$)
Camaná	72.73 °W; 16.60 °S	1969–2006	Pacific	Discharge ($\text{m}^3 \text{s}^{-1}$)
Huatiapa	72.47 °W; 16.00 °S	1969–2006	Pacific	Discharge ($\text{m}^3 \text{s}^{-1}$)
Titicaca	69.63 °W; 16.10 °S	1956–2014	Titicaca	Water level (m)
Rurrenabaque	67.54 °W; 14.44 °S	1968–2013	Amazon	Discharge ($\text{m}^3 \text{s}^{-1}$)

Data and method

Observed hydrological data

The low density of precipitation gauges in the Central Andes makes them generally inadequate for displaying spatio-temporal rainfall variability, which is largely influenced by complex orographic characteristics (e.g. Espinoza *et al* 2015). Discharge and water level data (hydrological dataset) are more reliable, because they represent accumulated precipitation in the catchment. Furthermore, the time period covered by the hydrological data (1956–2014) for the Central Andes is longer than that of meteorological data (generally starting after 1963; e.g. Espinoza *et al* (2009)). In this study we use a comprehensive hydrological dataset of discharge and water level data from the National Meteorology and Hydrology Service (SENAMHI) of Peru and the National Meteorology and Hydrology Service (SENAMHI) of Bolivia. The dataset covers a cross-section extending from the coast across the highlands of Peru and the Peruvian-Bolivian Altiplano to the Andean/Amazonian region of Bolivia. This dataset therefore provides a good indication of the different precipitation regimes in the Central Andes (15 °S–20 °S and 73 °W–63 °W; figures 1(a) and (b)).

For the Pacific system, we analyze the Huatiapa and Puente Camaná stations, in the Camaná River, as well as the La Tranca and Aguas Calientes stations in the Sama and Caplina Rivers, respectively. In the Tambo River, we use the La Pascana station (1956–1999) and the nearby Santa Rosa (1990–2014) station, which are highly correlated during the wet season ($r = 0.99$). We merge the data from these two stations to create one discharge time series for 1956 to 2014. In the Amazon drainage, discharge data from the Rurrenabaque station (1968–2013) on the Beni River is also analyzed. Discharge information for each station is calculated as the annual mean from December to April, which corresponds to rainfall during the wet season over this region. Finally, we use measurements of the Titicaca water level from December (year $n-1$) to April (year n) as an indicator of rainfall variability during the austral summer in the Altiplano (1914–2014). Table 1 displays the location, period and type of hydrological data for each station.

Oceanic and atmospheric data

Because our goal is to link the decadal–interdecadal hydrological variability of the Central Andes with oceanic patterns, we analyze sea surface temperature (SST; 1956–2014) data from Extended Reconstructed SST V3b (NOAA, Reynolds and Smith 1994). We also use zonal winds at 200 hPa (1966–2014) from the NCEP–NCAR reanalysis (Kalnay *et al* 1996) to analyze atmospheric processes associated with the decadal–interdecadal oscillation in the Central Andes.

We use the mean from December to April of the SST time series from 1956 to 2014 because that is our longest record of hydrological data. Zonal wind at 200 hPa (U200) over the Central Andes, however, shows a jump in the mean from 1949 to 1965, compared with the mean from 1949 to 2014 (which is not observed in other reanalysis data; figure S1). In addition, the Cramer test (Cramer 1946) shows that the difference in the means is significant at 90%. We therefore use only the mean from December to April of U200 time series from 1966 to 2014.

Defining the hydroclimatic index in the Central Andes

We standardized all hydrological data to compare eastern (wetter) and western (drier) regions and to compare discharge and water level data (figure 1(c), table 1). Each hydrological data is standardized subtracting its 50th percentile value and dividing by its standard deviation of each time series. To synthesize hydrological data (discharge and water level), we use the first principal component (PC1) resulting from an Empirical Orthogonal Function (EOF, Lorenz 1959), which is computed considering the standardized values of the seven hydrological data sets. Thus, the input matrix for the EOF is a 37×7 dimension (37 years for the common 1969 to 2005 period and seven standardized hydrological time series). We also propose a hydrological index over the Central Andes, which is computed as the mean of the standardized values of the discharge data for the Tambo River and the water level data for Titicaca Lake. We call this index Tam-Titi. To investigate the dominant frequency in our region of study, spectral analysis of PC1 and Tam-Titi is calculated using the multitaper power spectral density estimate (Thomson 1982). The red noise at 90% and 95% significance is computed to identify




Figure 2. (a) Correlation of the first principal component (PC1) with each hydrological station. Colors from white to dark red show the strength of the correlation. (b) Interannual variability of Titicaca-Tambo (Tam-Titi, black line) index, PC1 (blue line), and the low frequency of Tam-Titi (LTam-Titi; green). Wet (shading) and dry (no shading) periods are also shown. (c) Density power spectrum of Tam-Titi (black line) and red noise at 0.9 (blue) and 0.95 (red) significance.

leading frequencies. The decadal–interdecadal variability of Tam-Titi, SST and zonal wind at 200 hPa is extracted by applying a Butterworth low pass filter with a cutoff at 8 years and a scale of 2. Linear regressions and correlation are then performed between the low frequency variability of Tam-Titi and the SST (1957–2013) to identify oceanic zones that have an influence in the Central Andes. The same procedure is done with the zonal wind at 200 hPa (1967–2013).

Since correlation is performed between two filtered time series, which means that the autocorrelation of the time series could influence the correlations coefficient (Bertherton *et al* 1999), we calculate the significance of each correlation with modified degrees of freedom proposed by Bertherton *et al* (1999).

Determining the hydroclimatic low frequency variability in the Central Andes

The interannual hydrological variability of all stations correlates well (figure 1(c)), with a correlation coefficient that varies from 0.38; $p < 0.05$ (Aguas Caliente versus Rurrenabaque) to 0.91; $p < 0.01$ (Camana versus Huatiapa). Figure 1(c) shows a dominant oscillation of between 8 and 20 years, which is common in all the stations, explaining dry and wet periods over the Central Andes. This common oscillation suggests that a decadal–interdecadal low frequency is characteristic of the hydroclimate of the Central Andes.

To synthesize the main mode of hydroclimatic variability over the Central Andes, we conducted an Empirical Orthogonal Function (EOF) analysis for the seven stations. PC1 (68% of total variance) has a highly significant and positive correlation with all the hydroclimatic time series (figure 2(a)). The minimum correlation coefficient (r) is 0.71 (PC1 versus Rurrenabaque) and the maximum r is 0.94 (PC1 versus Tambo), both with a $p < 0.01$ (figure 2(a)). PC1 can therefore be considered an index that summarizes the variability of the hydrological stations. PC1 is also characterized by a low frequency with wet and dry periods that fluctuate between 8 and 20 years (blue line in figure 2(b)).

Although PC1 is a good indicator of variability in the Central Andes, its time period is relatively short (1969–2005) and is limited to the common period of the hydrological stations. To obtain an index covering a longer period (1956–2014), we computed the average of standardized values of discharge data for Tambo River and water level data for Lake Titicaca (Tam-Titi). Tam-Titi is well correlated to PC1 ($r = 0.93$, $p < 0.01$) for the common time period (1969–2005); it also shows decadal–interdecadal fluctuations of dry and wet hydrological conditions (black line in figure 2(b)). The correlation between Tam-Titi and the average of the remaining five stations is 0.85 ($p < 0.01$). These correlation coefficients suggest that Tam-Titi can be considered representative of hydrological variability in Central Andes.

The density power spectrum (DPS) of Tam-Titi and PC1 are calculated using the Thompson technique

(Thomson 1982). The largest peak in the DPS of Tam-Titi (figure 2(c)) and PC1 (figure S2) is between 8 to 20 years (frequency between 0.05 to 0.125 years⁻¹). This peak in the Tam-Titi DPS is significant at $p < 0.05$ (figure 2(c)), taking red noise into account, and the peak in the PC1 DPS is significant at $p < 0.10$ (figure S2). Those results are statistically robust enough to indicate that there is a marked decadal–interdecadal hydroclimatic variability in the Central Andes.

Decadal hydrological variability and ocean and atmospheric conditions

The decadal–interdecadal variability of hydrological time series in Central Andes could be a response to atmospheric and oceanic forcings. To evaluate these influences, we use the low frequency variability of Tam-Titi, which is calculated by retaining variability with periods greater than 8 years. This new time series (LTam-Titi) is marked by dry and wet conditions in the same periods as Tam-Titi (green line in figure 2(b)).

The central-western tropical Pacific Ocean explains 45% to 60% of the variability of LTam-Titi (figure 3(a)). This suggests that El Niño regions are not the main forcing of the hydrological variability of the Central Andes at a decadal–interdecadal time scale. The relationship between the low frequency of SST in the central-western tropical Pacific and LTam-Titi is negative, which means that when SST in the central-western tropical Pacific is cooler (warmer) than normal, LTam-Titi shows positive (negative) anomalies (figures 3(a) and (e)). In addition, climatic indices such as PDO, SST in the north and south tropical Atlantic and SST of the Niño 3.4 and Niño 1 + 2 regions are analyzed, but they do not show a robust relationship with LTam-Titi (table S1). These results suggest that the decadal variability described by LTam-Titi in the Central Andes do not necessarily agree with decadal rainfall variability previously described in South America, usually related to PDO (e.g. Marengo 2004, Andreoli and Kayano 2005, Garreaud *et al* 2008, Seiler *et al* 2013). Table S1 also shows a significant correlation between LTam-Titi and ENSO Modoki Index (EMI). EMI is computed using three zones of the tropical Pacific, including western, central and eastern equatorial Pacific Ocean (Ashok *et al* 2007). However, according to figure 3(a) significant correlation between low frequency SST and LTam-Titi only appears in the central-western tropical Pacific Ocean, suggesting a different spatial structure than EMI in the equatorial Pacific.

Low frequency of zonal wind at 200 hPa (U200) above the central Andes explains between 45% and 75% of the variance of LTam-Titi (figure 3(b)). The low frequency of the U200 is therefore related with LTam-Titi at a decadal–interdecadal time scale, and the relationship between them is negative (figures 3(b)

and (e)). During decades when the low frequency of U200 has negative (positive) anomalies, more (less) precipitation is expected, and high (low) values of LTam-Titi are observed. Decades with high (low) LTam-Titi values appear to be characterized by strong intensity of easterly (westerly) wind anomalies. These results are consistent with Garreaud *et al* (2003), which showed that U200 over the Central Andes plays an important role in the variability of precipitation at seasonal and interannual time scales in this region, since easterly anomalies carry moisture from the inner part of the continent. Nevertheless, our results reveal that decadal–interdecadal variability of U200 above the Central Andes also controls low frequency rainfall variability over this region.

A DPS analysis of SST over the central-western tropical Pacific and U200 over the Central Andes shows that both indices are dominated by decadal–interdecadal variability with significant values of between 8 and 14 years (frequency between 0.125 and 0.07 years⁻¹), which coincide with the Tam-Titi DPS (Figures 3(c) and (d)). Extracting the low frequency time series of SST over the central-western Pacific Ocean (180 °W–150 °W; 2.5 °S–15 °N) and U200 over the Central Andes (75 °W–66 °W; 20 °S–15 °S), and comparing with LTam-Titi, figure 3(e) shows that years with high (low) values of LTam-Titi are characterized by negative (positive) anomalies of both low frequency SST over the central-western Pacific Ocean and U200 over the Central Andes. Indeed, during cold (warm) periods of low frequency SST, normalized anomaly of low frequency U200 and LTam-Titi are -1.04 (0.94) and 0.78 (-0.83), respectively (table 2). These anomalies are significantly different comparing to the long-term mean values of low frequency U200 and LTam-Titi, according to a t-test at $p < 0.01$. This result suggests that cold (warm) periods of low frequency SST enhance easterlies (westerlies) anomalies, which significantly increase (decrease) the rainfall over the central Andes (table 2).

The relationship between SST in the central-western Pacific Ocean and the low frequency of U200 above the Andes is evaluated using a regression and correlation analysis (figure 4). The central-western Pacific Ocean explains 66% of the low frequency of U200 above the Andes and the relationship between them is positive; this means that during decades when the central-western Pacific Ocean is in a warm (cold) state, westerly (easterly) anomalies in low frequency variability of U200 are predominant above the Central Andes (figure 4).

In addition, figure 4 shows significant correlation between SST in the central-western Pacific Ocean and zonal winds at 200 hPa over the southern Africa, southeastern Asia and the northeastern Pacific Ocean. In coherence with these signals, previous papers have documented decadal rainfall variability over these regions related to SST in the tropical Pacific Ocean. Indeed, Zhang *et al* (2015) document decadal rainfall




Figure 3. Correlation maps of the low frequency variability of the Tambo-Titicaca index (Tam-Titi) with low frequency variability (periods > 8 years) of: (a) sea surface temperature (SST; 1956–2014) and (b) zonal wind at 200 hPa (U200; 1966–2014). Positive (solid lines) and negative (dashed lines) coefficients of Pearson correlation are shown in (a) and (b). Coefficient of determination (R^2) of grids, where correlation is significant at 95% ($p = 0.05$), are shaded in (a) and (b). (c) Density power spectrum (black line) of the central-western Pacific SST (180°W–150°W; 2.5°S–15°N; 1956–2014) and red noise (red dashed line) at 0.90 significance; (d) the same as (c), but for U200 over the Central Andes (75°W–66°W; 20°S–15°S; 1966–2014). (e) Standardized values of the low frequency variability (periods > 8 years) of the average of: SST (blue line) in the central-western Pacific, zonal winds at 200 hPa (blue line) over the Central Andes and Tam-Titi (black line). Note that SST and U200 are multiplied by -1 in (e).

Table 2. Periods of extreme warm (cold) low frequency sea surface temperature (SST) in the central west Pacific Ocean (180°W–150°W; 2.5°S–15°N), considering years of normalized SST values above (below) 0.5 (−0.5). Mean normalized anomalies for cold and warm periods of low frequency values of SST, zonal wind at 200 hPa (U200) over the central Andes (80°W–70°W; 19°S–14°S) and hydroclimatic index in the central Andes (LTam-Titi) are indicated.

Extreme SST low frequency conditions	Periods (years)	Normalized anomaly		
		SST	U200	LTam-Titi
Cold SST	1972–1977; 1998–2001; 2008–2012	−1.25	−1.04*	0.78*
Warm SST	1967–1970; 1980–1981; 1989–1996; 2003–2006	1.14	0.94*	−0.83*

*Values of U200 and LTam-Titi for warm and cold conditions are significantly different compared to their long-term mean according to t-test at $p < 0.01$.

variability over southern Africa modulated by SST in the central-western tropical Pacific. Zhu *et al* (2014) suggest decadal rainfall variability in southern China related to warming in the western Pacific Ocean and cooling in the eastern Pacific Ocean, which produces a strengthening of the Walker cell. Finally, Barlow *et al* (2001) point out that rainfall variability over south-

western USA at decadal time scale is associated with SST in the central-western tropical Pacific Ocean. These results might suggest the role of decadal variability of SST in the central-western Pacific Ocean in modulating rainfall in several regions; nevertheless the atmospheric processes related to these teleconnections are still not fully understood.




Figure 4. Correlation map of low frequency variability (periods >8 years) of SST (180 °W–150 °W; 2.5 °S–15 °N; 1966–2014), with low frequency variability of zonal wind at 200 hPa (U200). Positive (negative) values of correlation are drawn in solid (dotted) line. In addition, coefficient of determination (R^2 , shaded) of grids that have a correlation with significance greater than 95% ($p = 0.05$) is shown.

Discussion of the hydrological impacts of El Niño events

Extremely dry conditions occurred in the Central Andes during the 1982–83 El Niño ((Garreaud and Aceituno 2001; figures 1(c) and 2(b)); during the 1997–98 El Niño, however, which was one of the warmest El Niño events (MacPhaden 1999), the Central Andes did not show extremely dry conditions at all of the stations (figure 1(c)). Meanwhile, 1965, 1966, 1980, 1990, 1992 and 2005 were not classified as extreme El Niño years, but there were extremely dry conditions in Central Andes (figures 2(b) and 3(c)). These extreme impacts in the hydroclimate over this region can be explained by the low frequency modulation of LTam-Titi, since the years 1965, 1966, 1980, 1990, 1992 and 2005 fall within decades of lower LTam-Titi values (figures 2(b) and 3(e)). These results therefore suggest that low frequency variability plays a relevant role in modulating extreme hydrological impacts over the Central Andes.

Final comments

Analyzing annual hydrological data for the Central Andes and using density power spectral, we identified a significant decadal–interdecadal oscillation that controls the hydroclimatic variability of this region, which includes Peruvian and Bolivian Andes/Amazon and Altiplano and northern Chilean Andes. Following a regression analysis, our results suggest that the central-western tropical Pacific is the principal source of the decadal–interdecadal variability of the Central Andes, which influences low frequency variability of zonal wind at 200 hPa (U200) over this region. Decades characterized by

cold (warm) conditions in the central-western Pacific Ocean are associated with easterly (westerly) anomalies of U200 over the Andes. These easterly (westerly) anomalies enhance (inhibit) conditions for precipitation over the Central Andes at a decadal–interdecadal time scale. The results provided in this study indicate that while El Niño plays a relevant role in modulating rainfall variability over central Andes at interannual time scale, central-western Pacific SST have a major influence at decadal–interdecadal time scale. Finally, our results suggest that the intensity of hydrological impacts over the Central Andes during El Niño events might be influenced by the low frequency variability of central-western Pacific SST and U200 over this region.

Acknowledgments

We thank the following agencies/organizations for providing access to data: the National Meteorology and Hydrology National Service (SENAMHI) of Peru, the National Meteorology and Hydrology National Service (SENAMHI) of Bolivia, and the Observation Service for the geodynamical, hydrological and biogeochemical control of erosion/alteration and material transport in the Amazon, Orinoco and Congo basins (SO-HYBAM) for the hydrological data set used in this study; and NOAA/OAR/ESRL PSD, Boulder, Colorado, USA (www.esrl.noaa.gov/psd/) for the NOAA_ERSST_V4 sea surface temperature and NCEP reanalysis zonal wind data. We also thank PNICP-Peru for providing funds through contract 397-PNICP-PIAP-2014. We are grateful to Dr Kobi Mosquera, Dr Ivonne Montes, Ing. Yakelyn Ramos and Ing. Francisco Segura for their assistance with the

discussion of our results. We are grateful to Barbara Fraser for her contribution to improve this letter.

References

- Aceituno P and Garreaud R 1995 Impacto de los fenómenos el Niño y la Niña en regímenes fluviométricos andinos *Rev. Chilena de Ing. Hidráulica* **10** 33–43
- Andreoli R V and Kayano M T 2005 ENSO-related rainfall anomalies in South America and associated circulation features during warm and cold Pacific decadal oscillation regimes *Int. J. Climatol.* **25** 2017–30
- Ashok K, Behera S K, Rao S A, Weng H and Yamagata T 2007 El Niño Modoki and its possible teleconnection *J. Geophys. Res.* **112** C11007
- Barlow M, Nigam S and Wallace J M 2001 ENSO, Pacific decadal variability, and US summertime precipitation, drought, and stream flow *J. Clim.* **14** 2105–28
- Bretherton C S, Widmann M, Dymnikov V P, Wallace J M and Blade I 1999 The effective number of spatial degrees of freedom of a time-varying field *J. Clim.* **12** 1990–2009
- Cramer H 1946 *Mathematical Methods of Statistics* (Princeton: Princeton University Press) p 82
- Espinoza J C, Ronchail J, Guyot J L, Cochonneau G, Naziano F, Lavado W, De Oliveira E, Pombosa R and Vauchel P 2009 Spatio-temporal rainfall variability in the Amazon basin countries (Brazil, Peru, Bolivia, Colombia, and Ecuador) *Int. J. Climatol.* **29** 1574–94
- Espinoza J C, Chavez S, Ronchail J, Junquas C, Takahashi K and Lavado W 2015 Rainfall hotspots over the southern tropical Andes: spatial distribution, rainfall intensity, and relations with large-scale atmospheric circulation *Water Resour. Res.* **51** 3459–75
- Garreaud R 2000 Intraseasonal variability of moisture and rainfall over the South American Altiplano *Mon. Weather Rev.* **128** 3337–46
- Garreaud R, Vuille M, Compagnucci R and Marengo J 2008 Present-day South American climate *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **281** 180–95
- Garreaud R and Aceituno P 2001 Interannual rainfall variability over the South American Altiplano *J. Clim.* **14** 2779–89
- Garreaud R D 1999 Multiscale analysis of the summertime precipitation over the central Andes *Mon. Weather Rev.* **127** 901–21
- Garreaud R D 2009 The Andes climate and weather *Adv. Geosci.* **22** 3–11
- Garreaud R D, Vuille M and Clement A C 2003 The climate of the Altiplano: observed current conditions and mechanisms of past changes *Palaeoclimatology* **19** 45–22
- Houston J and Hartley A J 2003 The central Andean west-slope rainshadow and its potential contribution to the origin of hyper-aridity in the Atacama Desert *Int. J. Climatol.* **23** 1453–64
- Insel N, Poulsen C J and Ehlers T A 2010 Influence of the Andes mountains on South American moisture transport, convection, and precipitation *Clim. Dyn.* **35** 1477–92
- Kalnay E *et al* 1996 The NCEP/NCAR 40-year reanalysis project *Bull. Am. Meteorol. Soc.* **77** 437–71
- Knüsel S, Brüttsch S, Henderson K A, Palmer A S and Schwikowski M 2005 ENSO signals of the twentieth century in an ice core from Nevado Illimani, Bolivia *J. Geophys. Res.* **110** D01102
- Lagos P, Silva Y, Nickl E and Mosquera K 2008 El Niño-related precipitation variability in Peru *Adv. Geosci.* **14** 231–7
- Lavado-Casimiro W and Espinoza J C 2014 Impacts of El Niño and La Niña in the precipitation over Perú (1965–2007) *Revista Brasileira de Meteorologia* **29** 171–82
- Lenters J D and Cook K H 1997 On the origin of the Bolivian high and related circulation features of the South American climate *J. Atmos. Sci.* **54** 656–77
- Lorenz E 1959 Empirical orthogonal functions and statistical weather prediction *Scientific Report 1* (Department of Meteorology, Massachusetts Institute of Technology)
- Marengo J A 2004 Interdecadal variability and trends of rainfall across the Amazon basin *Theor. Appl. Climatol.* **78** 79–96
- McPhaden M 1999 Genesis and evolution of the 1997–98 El Niño *Science* **283** 950–4
- Melice J L and Roucou P 1998 Decadal time scale variability recorded in the Quelccaya summit ice core $\delta^{18}\text{O}$ isotopic ratio series and its relation with the sea surface temperature *Clim. Dyn.* **14** 117–32
- Minvielle M and Garreaud R D 2011 Projecting rainfall changes over the South American Altiplano *J. Clim.* **24** 4577–83
- Reynolds R W and Smith T M 1994 Improved global sea surface temperature analyses using optimum interpolation *J. Clim.* **7** 929–48
- Ronchail J 1995 Variabilidad interanual de las precipitaciones en Bolivia *Bull. Inst. Fr. Etud. Andines* **46** 13–33
- Rutllant J A, Fuenzalida H and Aceituno P 2003 Climate dynamics along the arid northern coast of Chile: the 1997–1998 Dinámica del Clima de la Región de Antofagasta (DCLIMA) experiment *J. Geophys. Res.* **108** 4538
- Seiler C, Hutjes R W and Kabat P 2013 Climate variability and trends in Bolivia *J. Appl. Meteorol. Clim.* **52** 130–46
- Silva Dias P L, Schubert W H and DeMaria M 1983 Large-scale response of the tropical atmosphere to transient convection *J. Atmos. Sci.* **40** 2689–707
- Sperling F, Valdivia C, Quiroz R, Valdivia R, Angulo L, Seimon A and Noble I 2008 Transitioning to climate resilient development: perspectives from communities in Peru *Environment Department #papers No. 115. Climate Change Series* (Washington, DC: World Bank) (<http://documents.worldbank.org/curated/en/619571468293754465/Transitioning-to-climate-resilient-development-perspectives-from-communities-in-Peru>)
- Thomson D J 1982 Spectrum estimation and harmonic analysis *Proc. IEEE* **70** 1055–96
- Vuille M 1999 Atmospheric circulation over the Bolivian Altiplano during dry and wet periods and extreme phases of the Southern Oscillation *Int. J. Climatol.* **19** 1579–600
- Vuille M, Bradley R S, Healy R, Werner M, Hardy D R, Thompson L G and Keimig F 2003 Modeling $\delta^{18}\text{O}$ in precipitation over the tropical Americas: 2. Simulation of the stable isotope signal in Andean ice cores *J. Geophys. Res.* **108** 4175
- Vuille M, Bradley R S and Keimig F T 2000 Interannual climate variability in the Central Andes and its relation to tropical Pacific and Atlantic forcing *J. Geophys. Res.: Atmos.* **105** 12447–60
- Vuille M and Keimig F 2004 Interannual variability of summertime convective cloudiness and precipitation in the central Andes derived from ISCCP-B3 data *J. Clim.* **17** 3334–48
- Zhang Q, Holmgren K and Sundqvist H 2015 Decadal Rainfall dipole oscillation over southern africa modulated by variation of austral summer land–sea contrast along the east coast of Africa *J. Atmos. Sci.* **72** 1827–36
- Zhu Z, Li T and He J 2014 Out-of-phase relationship between boreal spring and summer decadal rainfall changes in southern China *J. Clim.* **27** 1083–99